arXiv:1405.7790v2 [math.GT] 2 Jun 2014

Branched coverings of Seifert fibered solid tori *

Hong Huang

Abstract

In a paper published twelve years ago, the author gave a criterion to determine whether there is a fiber preserving branched covering between two given Seifert manifolds, which relies on the special case when the manifolds considered are Seifert fibered solid tori. Here we give an explicit construction of branched covering of a Seifert fibered solid torus by another when their Seifert invariants satisfy certain relation.

Key words: Seifert fibered solid torus, branched covering, Seifert invariant.

Mathematics Subject Classification 2010: 57N10

In a paper [1] published twelve years ago, the author gave a criterion to determine whether there is a fiber preserving branched covering between two given orientable Seifert manifolds with orientable bases, which relies on the special case when the manifolds considered are Seifert fibered solid tori. Here we give an explicit construction of fiber preserving branched covering of a Seifert fibered solid torus by another when their Seifert invariants satisfy certain relation.

Let V be a Seifert fibered solid torus. Let (q, h) be a (cross section, fiber) basis of $H_1(\partial V)$. We will abuse notation to use $q, h, etc.$ to denote both a closed curve and the homology class that it represents. If the meridian m of V is homologous to $\alpha q + \beta h$ in $H_1(\partial V)$, where $\alpha > 0, \beta$ are coprime integers, then we say that the fibered solid torus V has Seifert invariant (α, β) (w.r.t. (q, h)); see [2]. The following result is contained in [1], but the details of the proof of sufficiency are omitted there.

Proposition 0.1. *([1]) Let V_1, V_2 be two Seifert fibered solid tori. Suppose the Seifert invariant of V_2 w.r.t. some (cross section, fiber) basis of $H_1(\partial V_2)$ is (α_2, β_2), and k is a positive integer. Then there is a fiber preserving k-fold branched covering $f : V_1 \to V_2$ with fiber degree one and with upstairs and downstairs branched set the central fiber of V_1 and V_2 respectively if and only if there is some (cross section, fiber) basis of $H_1(\partial V_1)$ such that the Seifert invariant (α_1, β_1) of V_1 w.r.t. this basis satisfies $\beta_1/\alpha_1 = k\beta_2/\alpha_2$.*

*Research supported by NSFC (No. 11171025) and by Laboratory of Mathematics and Complex Systems, Ministry of Education.

Proof First we show the necessity as in [1]. Let $f : V_1 \to V_2$ be a fiber preserving k-fold branched covering with fiber degree one and with upstairs and downstairs branched set the central fiber of V_1 and V_2 respectively. (By upstairs branched set we mean the preimage of the downstairs branched set under f.) Let (q_2, h_2) be the (cross section, fiber) basis of $H_1(\partial V_2)$ w.r.t. which the Seifert invariant of V_2 is (α_2, β_2). Let $q_1 = f^{-1}(q_2)$. Since ∂V_1 is the union of fibers through all points of q_1 and the fiber degree of f is one, q_1 must be connected as ∂V_1 is. Then we see that q_1 is a cross section of the Seifert fibration of ∂V_1, and $f|_*(q_1) = kq_2$, where $f|$ is the restriction of f to ∂V_1.

Let the Seifert invariant of V_1 w.r.t. the (cross section, fiber) basis (q_1, h_1) be (α_1, β_1). By definition of branched covering and the assumption, there is a nonzero integer s such that $f|_*(m_1) = sm_2$, where m_1 (resp. m_2) is the meridian of V_1 (resp. V_2). So $f|_*(\alpha_1 q_1 + \beta_1 h_1) = s(\alpha_2 q_2 + \beta_2 h_2)$, and $\alpha_1 k q_2 + \beta_1 h_2 = s\alpha_2 q_2 + s\beta_2 h_2$, which implies that $\alpha_1 k = s\alpha_2$, and $\beta_1 = s\beta_2$. It follows that $\beta_1/\alpha_1 = k\beta_2/\alpha_2$.

Then we show the sufficiency. Suppose there is some (cross section, fiber) basis (q_1, h_1) of $H_1(\partial V_1)$ such that the Seifert invariant (α_1, β_1) of V_1 w.r.t. this basis satisfies $\beta_1/\alpha_1 = k\beta_2/\alpha_2$. If $\beta_2 = 0$, then $\alpha_2 = 1$, $\beta_1 = 0$, and $\alpha_1 = 1$. In this case, the result is clear. Below we assume that $\beta_2 \neq 0$. From the relation $\beta_1/\alpha_1 = k\beta_2/\alpha_2$, using that α_j and β_j are coprime, we get that $\alpha_1 | \alpha_2$, and $\beta_2 | \beta_1$.

Now it is clear that there exists a fiber preserving k-fold branched covering $f : V_1 \to V_2$ with fiber degree one and with upstairs and downstairs branched set the central fiber of V_1 and V_2 respectively. The reason is as follows. First we can construct a fiber preserving covering $f| : \partial V_1 \to \partial V_2$ such that $f|_*(q_1) = kq_2$, and $f|_*(h_1) = h_2$. We have, using the relation $\beta_1/\alpha_1 = k\beta_2/\alpha_2$,

$$f|_*(m_1) = f|_*(\alpha_1 q_1 + \beta_1 h_1) = \alpha_1 k q_2 + \beta_1 h_2 = \frac{\beta_1}{\beta_2}(\alpha_2 q_2 + \beta_2 h_2) = \frac{\beta_1}{\beta_2} m_2.$$

Then we shrink the tori ∂V_1 and ∂V_2 gradually (towards the central fibers) , and modify the map $f|$ correspondingly. At last we will get the desired branched covering $f : V_1 \to V_2$.

Below we give a more explicit construction. Let α_j', β_j' be integers such that $\alpha_j \beta_j' - \alpha_j' \beta_j = 1$, $j = 1, 2$; note that α_j' (resp. β_j') is determined by α_j and β_j up to multiples of α_j (resp. β_j). Thus we have $\alpha_1 \beta_1' - (\frac{\beta_1}{\beta_2}\alpha_1')\beta_2 = 1$, and $\alpha_1(\frac{\alpha_2}{\alpha_1}\beta_2') - \alpha_2'\beta_2 = 1$. We see that $\alpha_2' - \frac{\beta_1}{\beta_2}\alpha_1'$ is a multiple of α_1.

For $j = 1, 2$, consider the universal covering $p_j : D^2 \times \mathbb{R} \to V_j$ with deck transformation group generated by the element

$$\tau_j : D^2 \times \mathbb{R} \to D^2 \times \mathbb{R},$$
$$(re^{2\pi i\theta}, t) \mapsto (re^{2\pi i(\theta - \frac{\alpha_j'}{\alpha_j})}, t + 1).$$

Here we think of V_j as the quotient of $D^2 \times \mathbb{R}$ by the action of the corresponding deck transformation group; cf. [2].

First we construct a map $\tilde{f} : D^2 \times \mathbb{R} \to D^2 \times \mathbb{R}$ as follows,

$$\tilde{f}(re^{2\pi i\theta}, t) = (re^{2\pi i\frac{\beta_1}{\beta_2}\theta}, \frac{\alpha_2}{\alpha_1}t).$$

Note that

$$\begin{aligned}
&\tilde{f}(\tau_1(re^{2\pi i\theta}, t)) \\
&= \tilde{f}(re^{2\pi i(\theta-\frac{\alpha_1'}{\alpha_1})}, t+1) \\
&= (re^{2\pi i\frac{\beta_1}{\beta_2}(\theta-\frac{\alpha_1'}{\alpha_1})}, \frac{\alpha_2}{\alpha_1}(t+1)) \\
&= (re^{2\pi i(\frac{\beta_1}{\beta_2}\theta-\frac{\alpha_2}{\alpha_1}\frac{\alpha_2'}{\alpha_2})}, \frac{\alpha_2}{\alpha_1}t+\frac{\alpha_2}{\alpha_1}) \\
&= \tau_2^{\alpha_2/\alpha_1}(re^{2\pi i\frac{\beta_1}{\beta_2}\theta}, \frac{\alpha_2}{\alpha_1}t) \\
&= \tau_2^{\alpha_2/\alpha_1}(\tilde{f}(re^{2\pi i\theta}, t)).
\end{aligned}$$

So $\tilde{f}$ descends to a map $f : V_1 \to V_2$. One can easily check that $f : V_1 \to V_2$ is the desired branched covering. □

References

[1] H. Huang, Branched coverings and nonzero degree maps between Seifert manifolds, Proc. Amer. Math. Soc. 130 (2002), no. 8, 2443-2449.

[2] H. Seifert, Topologie dreidimensionaler gefaserter Raume, Acta Math. 60 (1933), 147-238. (There is an English translation by W. Heil.)

School of Mathematical Sciences, Beijing Normal University,
Laboratory of Mathematics and Complex Systems, Ministry of Education,
Beijing 100875, P.R. China
E-mail address: hhuang@bnu.edu.cn